

High-CO₂ Response Mechanisms in Microalgae

Masato Baba^{1,2} and Yoshihiro Shiraiwa^{1,2}

¹Graduate School of Life and Environmental Sciences,
University of Tsukuba, Tsukuba, Ibaraki,

²CREST, JST,
Japan

1. Introduction

The concentrations of atmospheric CO₂ and aquatic inorganic carbon have decreased over geologic time with minor fluctuations. In contrast, O₂ concentration has increased through the actions of photosynthetic organisms. Therefore, photosynthetic organisms must adapt to such dramatic environmental change. Aquatic photosynthetic microorganisms, namely eukaryotic microalgae, cyanobacteria, and non-oxygen-evolving photosynthetic bacteria, have developed the ability to utilize CO₂ efficiently for photosynthesis because CO₂ is a substrate for the primary CO₂-fixing enzyme ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) and its related metabolic pathways such as the Calvin-Benson cycle (C₃ cycle). As the Rubisco carboxylase reaction is suppressed by elevated O₂ concentrations via competition with CO₂, photosynthetic organisms have developed special mechanisms for acclimating and adapting to changes in both CO₂ and O₂ concentrations. Examples of such mechanisms are the microalgal CO₂-concentrating mechanisms (CCM), the facilitation of “indirect CO₂ supply” with the aid of carbonic anhydrase and dissolved inorganic carbon (DIC)-transporters (see Section 3), and C₄-photosynthesis (for review, see Giordano et al., 2005; Raven, 2010). Many reports on low-CO₂-acclimation/adaptation mechanisms have been published, particularly in relation to certain cyanobacteria and unicellular eukaryotes. However, knowledge of high-CO₂-acclimation/adaptation mechanisms is very limited. We recently identified an acceptable high-CO₂-inducible extracellular marker protein, H43/Fea1 (Hanawa et al., 2007) and a *cis*-element involved in high-CO₂-inducible gene expression in the unicellular green alga *Chlamydomonas reinhardtii* (Baba et al., 2011a). We also identified other high-CO₂-inducible proteins in the same alga using proteomic analysis (Baba et al. 2011b). In this chapter, we briefly introduce low-CO₂-inducible phenomena and mechanisms as background and then review recent progress in elucidating the molecular mechanisms of the high-CO₂ response in microalgae.

2. Aquatic inorganic carbon system

The CO₂ concentration dissolved in aqueous solution (dCO₂) is equilibrated with the partial pressure of atmospheric CO₂ (pCO₂) by Henry’s law and depends on various environmental factors such as temperature, Ca²⁺ and Mg²⁺ levels, and salinity (e.g., Falkowski & Raven,

2007). The $d\text{CO}_2$ dissociates into bicarbonate (HCO_3^-), and carbonate (CO_3^{2-}) and these three species of DIC attain equilibrium at a certain ratio depending on pH, ion concentrations, and salinity (Fig. 1). HCO_3^- is the dominant species at physiological pH (around 8), which is similar to that in the chloroplast stroma where photosynthetic CO_2 fixation is actively driven (for review, see Bartlett et al., 2007). However, Rubisco [E.C. 4.1.1.39] reacts only with $d\text{CO}_2$, not bicarbonate or carbonate ions. At a pH of 8, the $d\text{CO}_2/\text{HCO}_3^-$ ratio becomes extremely small (approximately 1/100) resulting in a high bicarbonate concentration and an increase in the total DIC pool size. The $d\text{CO}_2$ concentration equilibrates with atmospheric CO_2 at approximately 10 μM , whereas the bicarbonate concentration is approximately 2 mM at the surface of the ocean (Falkowski & Raven, 2007).




Fig. 1. Equilibration of dissolved inorganic carbon species in freshwater and seawater. Parameters used were as follows (at 25°C): For freshwater, $pK_{a1} = 6.35$, $pK_{a2} = 10.33$; for seawater, $pK_{a1} = 6.00$, $pK_{a2} = 9.10$ (Table 5.2, Falkowski & Raven, 2007). Filled symbols and solid line, freshwater; clear symbols and dotted line, seawater; diamonds, $d\text{CO}_2$; squares, bicarbonate; triangles, carbonate.

CO_2 must be supplied rapidly when it is actively fixed by Rubisco in the chloroplast stroma during photosynthesis. CO_2 is supplied by both diffusion from outside of cells and the conversion of bicarbonate. However, these processes are very slow and become limiting for photosynthetic CO_2 fixation. In the former case, CO_2 must be continuously transported from outside of the cells via the cytoplasm through the plasmalemma and the chloroplast envelope. The diffusion rate of CO_2 in water is approximately 10,000-fold lower than that in the atmosphere (Jones, 1992). In the latter case, bicarbonate accumulated in the stroma can be a substrate when the dehydration rate to convert bicarbonate to CO_2 is comparable to Rubisco activity. However, the rate of chemical equilibration between CO_2 and the bicarbonate ion is very slow relative to photosynthetic consumption of CO_2 (Badger & Price, 1994; Raven, 2001); the first-order rate constants of hydration (CO_2 to bicarbonate) and dehydration (bicarbonate to CO_2) are 0.025–0.04 s^{-1} and 10–20 s^{-1} , respectively, at 25°C (Ishii et al., 2000). Such CO_2 -limiting stress becomes a motive for photosynthetic organisms to develop unique CO_2 -response mechanisms.

3. The CO₂-concentrating mechanism and phenomena induced by CO₂ limitation

The atmospheric CO₂ level has gradually decreased over recent geological time with some fluctuations (Condie & Sloan, 1998; Falkowski & Raven, 2007; Giordano et al., 2005; Inoue, 2007), although it has been increasing rapidly due to CO₂ emissions from fossil fuels since the industrial revolution. Thus, photosynthetic organisms have adapted to utilize CO₂ efficiently for photosynthesis. Generally, eukaryotic microalgae and cyanobacteria have developed efficient CO₂-utilization mechanisms and exhibit high photosynthetic affinity for CO₂ when grown under CO₂-limiting conditions. Under elevated CO₂ conditions, they exhibit low affinity for CO₂, as enough CO₂ is available for photosynthesis. These properties can change over hours when photosynthetic microorganisms are grown under various CO₂ conditions (for review, see Miyachi et al., 2003) (Fig. 2).




Fig. 2. Relationship between photosynthetic rate and external dissolved inorganic carbon (DIC) concentration in microalgae grown under low-, high-, and extremely high-CO₂ conditions. A, low-CO₂-acclimated cells (grown in air with 0.04% CO₂); B, low-CO₂-acclimated cells treated with a carbonic anhydrase inhibitor (e.g., *Chlorella*); C, high-CO₂-acclimated cells (grown in air containing 1–5% CO₂) (e.g., *Chlamydomonas*); D, high-CO₂-acclimated cells (e.g., *Chlorella*); E, extremely high-CO₂-acclimated cells grown under >40% CO₂ conditions. This figure is modified from Miyachi et al. (2003).

High photosynthetic activity of low-CO₂-grown cells under a CO₂-limiting concentration is due to the CCM, which is induced by cellular acclimation to limiting CO₂ (e.g., Aizawa and Miyachi, 1986). Two main factors are essential for CCM: inorganic carbon transporters that facilitate DIC membrane transport of CO₂ and/or bicarbonate through the plasmalemma and the chloroplast envelope, and carbonic anhydrases (CAs), which facilitate diffusion by stimulating the indirect supply of CO₂ from outside of cells to Rubisco. CA catalyzes the equilibration reaction of the hydration and dehydration of CO₂ and bicarbonate, respectively. The rapid equilibration catalyzed by CA stimulates the increase in bicarbonate concentration at physiological pH and augments the contribution of bicarbonate for diffusion. Finally, the processes driven by CA induce increases in the amount of bicarbonate

carried near Rubisco and then CO₂ produced from bicarbonate is immediately supplied to Rubisco when CA is located near Rubisco (see also Fig. 4). The relative specificity to CO₂/O₂ and affinity to CO₂ of Rubisco became more efficient over evolutionary time, indicating that Rubisco in eukaryotic microalgae is more efficient for CO₂ fixation than that in cyanobacteria (Falkowski & Raven, 2007). Such species-specific properties remain unchanged in present living organisms. However, even in eukaryotic algae, the affinity of Rubisco for CO₂ is insufficient to saturate activity at present atmospheric CO₂ concentrations. Therefore, the cells continuously activate mechanisms such as CCM to increase their affinity for CO₂. For further information on CCM, we recommend reading several previously published reviews (e.g., Aizawa & Miyachi, 1986; Badger et al., 2006; Giordano et al., 2005; Kaplan & Reinhold, 1999; Miyachi et al., 2003; Moroney & Ynalvez, 2007; Raven et al., 2008; Raven, 2010; Spalding, 2008; Yamano & Fukuzawa, 2009).

CCM is reversibly induced/suppressed by the decrease/increase in CO₂ concentration, respectively, in cyanobacteria and eukaryotic microalgae when the duration of acclimation is on an hour- or day-order length. However, in the unicellular green alga *Chlamydomonas reinhardtii* in which CCM is the most characterized among eukaryotic microalgae, cells grown for 1000 generations under high-CO₂ conditions are unable to re-acclimate to low-CO₂ conditions, exhibiting low photosynthetic affinity for CO₂ even when the cells are re-exposed to low CO₂ conditions (Collins & Bell, 2004; Collins et al., 2006). This suggests that CCM can be irreversibly lost when cells undergo prolonged acclimation/adaptation to high-CO₂ conditions. Such adaptation has been suggested to occur in natural populations (Collins & Bell, 2006). Although CCM-deficient mutants of cyanobacteria and the green alga *C. reinhardtii* are lethal, such lethality is prevented by elevated CO₂ concentration (e.g., Price & Badger, 1989; Spalding et al., 1983; Suzuki & Spalding, 1989). In *C. reinhardtii*, CCM is induced under either 1.2% CO₂ in air at 1000 μmol photons m⁻² s⁻¹ or 0.04% CO₂ in air at 120 μmol photons m⁻² s⁻¹, suggesting that CCM induction can be regulated by not only external CO₂ concentration but also other signals derived from photorespiratory and/or excess photoenergy stresses, although the detailed mechanisms are not yet known (Yamano et al., 2008). CCM can be induced by an artificially produced strong limitation in CO₂ supply in large-scale photobioreactors where dCO₂ is consumed via photosynthesis (Yun & Park, 1997).

In some microalgae, the supply of CO₂, not bicarbonate, is strongly limited at alkaline pHs in closed culture systems and such a limitation may be a factor or signal for inducing CCM (Colman & Balkos, 2005; Diaz & Maberly, 2009; Verma et al., 2009). The euglenophyte *Euglena mutabilis* and an acid-tolerant strain of *Chlamydomonas* do not induce CCM under any conditions (Balkos & Colman, 2007; Colman & Balkos, 2005), suggesting that photosynthetic carbon fixation is not limited by CO₂ supply even under ambient atmospheric conditions. These results indicate that there is species-specific variation in the induction mechanism of CCM depending on physiological and ecological conditions (for review, see Giordano et al., 2005; Raven, 2010).

4. High-CO₂ response phenomena

The atmospheric CO₂ level is presumed to have been very high during the ancient geological era (Condie & Sloan, 1998; Falkowski & Raven, 2007; Giordano et al., 2005; Inoue, 2007), so microalgae are believed to have been high-CO₂-adapted/acclimated cells. Microalgae preserve their ancient physiological properties at present, and the relative

specificity of Rubisco is a typical example. Even in the present environment, high-CO₂ conditions occur in soil where CO₂ concentration changes drastically between the atmospheric level and $\geq 10\%$ (v/v) (for review, see Buyanovsky & Wagner, 1983; Stolzy, 1974). Accordingly, phenomena that are induced under high-CO₂ conditions, such as high-CO₂ acclimation, remain important for microalgae to survive in various environments.

Among the various phenomena induced by high CO₂ concentrations, keenly interesting topics are how to maximize inorganic CO₂ fixation and organic production by microalgae for CO₂ mitigation and mass cultivation. The most frequently used species for studies on fast growth and tolerance to high CO₂ levels is *Chlorella sp.*, followed by *Scenedesmus sp.*, *Nannochloropsis sp.*, and *Chlorococcum sp.* The CO₂ concentration used for such studies varies from atmospheric levels to 100% (Kurano et al., 1995; Maeda et al., 1995; Olaizola, 2003; Seckbach et al., 1970). Appropriate CO₂ supply for saturation of microalgal growth is approximately 5% in the unicellular green alga *Chlorella* (Nielsen 1955). The growth of microalgae and cyanobacteria is generally inhibited under very high concentrations of CO₂. Some species isolated from extreme environments can grow rapidly with tolerance to very high and extremely high CO₂ conditions such as >40% (for review, see Miyachi et al., 2003). Even in a high-CO₂-tolerant microalga, growth is suppressed at > 60% CO₂ in air (Satoh et al., 2004). The rate of maximum photosynthesis per packed cell volume increases in some species, such as *Chlorella*, but not in other species, such as *Chlamydomonas*, even when cells are acclimated to high-CO₂ conditions (Miyachi et al., 2003) (Fig. 2). However, the detailed mechanism on such high CO₂ tolerance needs to be clarified.

Many reports have focused on lipid biosynthesis for biofuel production, and response surface methodology (Box & Wilson, 1951) has been used very effectively to evaluate multiple factors associated with total biomass production. Excellent review articles on large-scale cultivation for biofuel production by microalgae and cyanobacteria have focused on how to obtain the best productivity under high-CO₂ conditions (Ho et al., 2011; Kumar et al., 2010; Lee J.S. & Lee J.P., 2003), but not on the underlying mechanisms of how cells provide high productivity under fine regulation.

One of the best examples of sequential analysis was performed systematically in the high-CO₂-tolerant unicellular green alga *Chlorococcum littorale* (for review, see Miyachi et al., 2003). *C. littorale* is a unicellular marine chlorophyte that was isolated from a saline pond in Kamaishi City, Japan; it grows rapidly under extremely high CO₂ conditions (e.g., 40%, and even at 60% CO₂; Chihara et al., 1994; Kodama et al., 1993; Satoh et al., 2004). Several experiments have revealed that cellular responses, namely the regulation of photosystem (PS) I and PS II, the production of ATP, and pH homeostasis are well maintained particularly in *C. littorale*, but not in high-CO₂-sensitive species such as the green soil alga *Stichococcus bacillaris*, during a lag period when cells are transferred from low to extremely high levels of CO₂ (Demidov et al., 2000; Iwasaki et al., 1996, 1998; Pescheva et al., 1994; Pronina et al., 1993; Sasaki et al., 1999; Satoh et al., 2001, 2002). However, many of the processes that make it possible for cells to grow under such extremely high-CO₂ conditions remain to be understood.

Photosynthesis in acidic environment, the influence by ocean acidification, and the effect of O₂ on photorespiration are also deeply associated with high-CO₂-induced phenomena. Some microalgal species have been isolated mainly from acidic environments where only CO₂ is predominant and supplied to algal cells as a substrate for photosynthesis (Balkos & Colman, 2007; Colman & Balkos, 2005; Diaz & Maberly, 2009; Verma et al., 2009; for review see Raven, 2010). Three synurophyte algae, *Synura petersenii*, *Synura uvella*, and *Tessellaria*

volvocina, have been studied in detail for the DIC uptake mechanism and show unique photosynthetic properties (Bhatti & Coleman, 2008). These species have no external carbonic anhydrase on the cell surface, no bicarbonate uptake ability, and exhibit a low affinity for DIC during photosynthesis, indicating a lack of CCM as in high-CO₂-grown/acclimated cells. However, their Rubisco shows a relatively high affinity for CO₂, and cells such as *S. petersenii* accumulate large amounts of internal DIC via diffusive uptake of CO₂ facilitated by a pH gradient across the cell membranes, as reported previously in spinach chloroplasts (Heldt et al., 1973). These data suggest that the affinity of Rubisco for CO₂ and the homeostasis of the pH gradient play key roles in the whole-cell affinity for CO₂ and the pH-tolerance of microalgae. Under high-CO₂ conditions, Rubisco can get enough CO₂ supply although CCM is usually lost in high-CO₂ cells. The physiological status of synurophyte algae living at acidic pH may be similar to cells that are exposed to high-CO₂ conditions even under low-CO₂ conditions.

Increasing pCO₂ induces a decrease in oceanic pH and causes gradual equilibrium shifts from bicarbonate ions to CO₂ in seawater. Therefore, ocean acidification is said to be another high-CO₂ problem (for review, see Doney et al., 2008). Coccolithophorids, marine phytoplankton that form cells covered with CaCO₃, are very sensitive to calcium carbonate saturation and pH shifts in seawater. The effects of ocean acidification on algal physiology have been studied in several coccolithophorid species such as *Emiliania huxleyi* and *Pleurochrysis carterae*, although some conflicting results have been reported (Fukuda et al., 2011; Igresiaz-Rhodorigez et al., 2008; Riebesell et al., 2000). Hurd et al. (2009) indicated the importance of maintaining pH in experiments and demonstrated that doing so via high-CO₂ bubbling creates conditions that are much closer to actual ocean acidification than acidification by adding HCl. The effects of high-CO₂ conditions on calcification and photosynthesis would be closed up in later analyses. Fukuda et al. (2011) reported that the coccolithophorid *E. huxleyi* possesses alkalization activity, which helps compensate for acidification when photosynthesis is actively driven. Furthermore, when oceanic acidification is caused by the bubbling of air with elevated CO₂, coccolithophorid cells increase both photosynthetic activity and growth and are not damaged because of the stimulation of photosynthesis (unpublished data by S. Fukuda, Y. Suzuki & Y. Shiraiwa). These results suggest that ocean acidification will not immediately harm coccolithophorids. However, long-term experimental evidence is strongly required on this topic.

Badger et al. (2000) described how low-CO₂-grown microalgae tend to have low photorespiratory activity, as determined by photosynthetic O₂ uptake in C₄ plants because of the function of CCM. O₂ uptake under illumination is relatively insensitive to changes in CO₂ concentration, because the activity depends predominantly on the activity of non-photorespiratory reactions probably such as the Mehler reaction and oxidizing reaction in the mitochondria (Badger et al., 2000). CO₂ insensitivity is also observed in *C. reinhardtii* (Sültemeyer et al., 1987) although photosynthetic O₂ uptake increases considerably with increasing light intensity (Sültemeyer et al., 1986). Accordingly, the photosynthetic productivity of microalgae may not be significantly enhanced by suppressing photorespiration. The rate of maximum photosynthesis, calculated on a cell volume, increases clearly in *Chlorella* but not so in *Chlamydomonas* when cells are acclimated to high-CO₂ conditions (Miyachi et al., 2003) (Fig. 2). In *C. reinhardtii*, growth rate is only slightly higher (1.3–1.8-fold) in cells grown under high-CO₂ than in those grown under ordinary air (Baba et al., 2011b; Hanawa, 2007). These results suggest that low-CO₂-acclimated/grown cells have a very highly efficient carbon-fixation mechanism for maintaining high growth

rates even under atmospheric CO₂ levels, so we need to carefully optimize growth conditions when we want to obtain high algal growth and production using CO₂ enrichment (see also section 5).

5. Molecular mechanisms for high-CO₂ responses

Microalgae can acclimate to high-CO₂ conditions by changing their photosynthetic properties such as CCM. The half-saturation concentration of CO₂ for changing cellular photosynthetic characteristics, i.e., CO₂ affinity, is 0.5% in the unicellular green alga *Chlorella kessleri* 211-11h (formerly *C. vulgaris* 11h; Shiraiwa & Miyachi, 1985). CCM-related proteins are also degraded simultaneously when cells are transferred from low- to high-CO₂ conditions (see references in section 3). Yang et al. (1985) found that, during acclimation to high-CO₂ conditions, CA, an essential component of CCM, was passively degraded and thus the process took almost 1 week.

C. reinhardtii cells in freshwater and in soil are exposed to drastically fluctuating concentrations of CO₂ between atmospheric level and $\geq 10\%$ (v/v) (for review, see Stolzy, 1974; Buyanovsky & Wagner, 1983). To grow in such habitats and maintain optimum growth, the alga needs to rapidly change its physiology. Such rapid acclimation was in fact observed in *C. reinhardtii* cells that were successfully acclimated to 20% CO₂ within a few days (Hanawa, 2007). The specific growth rate (μ) of *C. reinhardtii* was 0.176 in ordinary air containing 0.04% CO₂ where dCO₂ and total DIC were 1.62 and 6.19 μM , respectively, at pH 6.8 (Hanawa, 2007) (Fig. 3). Although dCO₂ and total DIC concentrations in the culture media, which were equilibrated with 0.3, 1.0, and 3.0% CO₂ (v/v) in air, were 28-, 121-, and 489-fold higher than that in ordinary air, respectively, alga-specific growth rates under the respective conditions were only 1.3-, 1.8-, and 1.7-fold higher than that in air (Hanawa, 2007) (Fig. 3). In a wall-less mutant of *C. reinhardtii* CC-400 (same as CW-15), the growth rate and the amount of total proteins increased only 1.5-fold even when the CO₂ concentration was increased from atmospheric level to 3% (Baba et al., 2011b). These results clearly indicate that, in *C. reinhardtii*, CO₂ enrichment is not advantageous to increase in growth rate, as the fully low-CO₂-acclimated cells acquire CCM and grow quickly with a near-maximum growth rate even under atmospheric levels of CO₂. These results are true when cells are growing logarithmically at low cell density to prevent self-shading. However, when cell density is quite high, the ratio of growth at high to low CO₂ is usually quite high. This is probably due to the decrease in growth under air conditions. Under such conditions, CO₂ supply is strongly limited resulting in very low growth rates under air-level CO₂. Nevertheless, the growth rate does not exceed the specific growth rate obtained at the logarithmic growth stage.

High-CO₂-grown *C. reinhardtii* declines CCM physiologically by losing CA and active DIC transport systems in order to avoid secondary inhibitory effects caused by excess DIC accumulation (for review, see Miyachi et al., 2003; Spalding, 2008; Yamano & Fukuzawa, 2009) but no other significant responses have been reported until recently. Recently, we found drastic changes in extracellular protein composition (Baba et al., 2011b) including induction of the H43/Fea1 protein (Hanawa et al., 2004, 2007; Kobayashi et al., 1997).

The wall-less mutant of *C. reinhardtii*, CW-15, releases a large amount of extracellular matrix, including periplasm-locating proteins, named as extracellular proteins, into the medium (Hanawa et al., 2007; Baba et al., 2011b). Our previous studies clearly showed that the extracellular protein composition changes drastically when *C. reinhardtii* cells are transferred




Fig. 3. Relationship between the specific growth rate and dCO₂ concentration in an air-bubbled culture of *Chlamydomonas reinhardtii* (A), and the concentrations of three dissolve inorganic carbon (DIC) species in the culture (B). The concentration of dCO₂ was experimentally determined. Each DIC species was calculated by Henley's law and the Henderson-Hasselbalch equation, respectively. The parameters were as follows (for freshwater at 25°C): pK_{a1} = 6.35, pK_{a2} = 10.33. The culture medium used was a high salt medium supplemented with 30 mM MOPS (pH 6.8). Crosses, specific growth rate; diamonds, dCO₂; circles, total DIC; squares, bicarbonate; triangles, carbonate. Fig. 3A is modified from Hanawa, 2007.

from atmospheric air to 3% CO₂ in air (Hanawa et al., 2004, 2007; Kobayashi et al., 1997), whereas an SDS-PAGE profile of intracellular-soluble and -insoluble proteins showed no clear difference (Baba et al., 2011b). Recently, we analyzed 129 proteins by proteomic analysis and identified 22 high-CO₂-inducible proteins from *C. reinhardtii* cells transferred from low- to high-CO₂ conditions (Baba et al., 2011b). These high-CO₂-inducible proteins are multiple extracellular hydroxyproline-rich glycoproteins (HRGPs), such as nitrogen-starved gametogenesis (NSG) protein (Abe et al., 2004), inversion-specific glycoprotein (ISG) (Ertl et al., 1992), and cell wall glycoprotein (GP) (Goodenough et al., 1986), together with sexual pherophorin (PHC) (Hallmann, 2006), gamete-specific (GAS) protein (Hoffmann & Beck, 2005), and gamete-lytic enzymes (Buchanan & Snell, 1988; Kinoshita et al., 1992; Kubo et al., 2001). Both GP and ISG are classified as HRGPs together with PHC, GAS, and sexual agglutinin with a shared origin (Adair, 1985). HRGPs are generally involved in sexual recognition of mating-type, plus or minus gametes, in the *Chlamydomonas* lineage (Lee et al., 2007). Among these proteins, NSG, GAS, and gamete-lytic enzymes are generally known to be induced during the gametogenetic process. The sexual program, including gametogenesis in *Chlamydomonas*, is strictly regulated by nitrogen availability (for review, see Goodenough et al., 2007). Drastic changes in the expression of gametogenesis-related extracellular proteins were clearly observed in *C. reinhardtii* cells in response to high-CO₂ but not to environmental nitrogen concentrations, because the experiment was performed under nitrogen-sufficient conditions (Baba et al., 2011b). No visible effect of high-CO₂ signal alone was observed on mating (Baba et al., 2011b). From these results, we concluded that the high-CO₂ signal induced gametogenesis-related proteins but that the signal was not strong

enough or was still missing some necessary factors to trigger mating. Otherwise, these gametogenesis-related protein families and/or HRGPs may have another function under high-CO₂ conditions.

The biological meaning of the expression of gametogenesis-related proteins at the stage of vegetative growth is quite mysterious. CCM may be differentially regulated by changes in nitrogen availability, depending on the species (for review, see Giordano et al., 2005). In *C. reinhardtii*, mildly limited nitrogen availability suppresses CCM and mitochondrial β -CA expression (Giordano et al., 2003) and the increase in NH₄⁺ concentration promotes the efficiency of photosynthetic CO₂ utilization (Beardall & Giordano, 2002). From these results, Giordano et al. (2005) suggested that the induction of CCM and related phenomena induced by CO₂ limitation is regulated to satisfy an adequate C/N ratio. Basically, cells growing under high-CO₂ conditions may require more nitrogen, at least no less than low-CO₂-acclimated cells, and tend to attain nitrogen-limitation status easily. In contrast, Giordano et al. (2005) suggested that activating CCM may reduce the loss of nitrogen through the photorespiratory nitrogen cycle. Namely, NH₄⁺ produced by converting Gly to Ser through the C₂ cycle in mitochondria is transported to and re-fixed in the chloroplasts by the GS2/GOGAT cycle where chloroplastic GS2 is induced in response to CO₂ concentration in *C. reinhardtii* (Ramazanov & Cárdenas, 1994). In previous works, the NH₄⁺ excretion rate from algal cells was lower in high-CO₂ cells than in low-CO₂ cells when monitored in the presence of 1 mM 1-methionine sulfoximine, a specific inhibitor of GS activity, to prevent re-fixation of NH₄⁺ in *C. reinhardtii* CW-15 (Ramazanov & Cárdenas, 1994) and similarly in *C. vulgaris* 211-11h (Shiraiwa & Schmid, 1986). A decrease in the intracellular NH₄⁺ level was first reported to induce gametogenesis-related genes in *C. reinhardtii* (Matsuda et al., 1992). Thus, it is reasonable to hypothesize that gametogenesis is triggered by a decrease in intracellular NH₄⁺ levels under high-CO₂ conditions when photorespiration is suppressed. However, further study is required, as photorespiratory activity in *C. reinhardtii* is very low (Badger et al., 2000).

Another report suggested the close participation of CO₂ in inorganic nitrogen assimilation (for review, see Fernández et al., 2009). LCIA, or NAR1.2, is involved in the bicarbonate transport system in chloroplasts (Duanmu et al., 2009) but is not regulated by nitrogen availability, and has been identified as a low-CO₂-inducible gene by expressed sequence tag (EST) analysis (Miura et al., 2004). However, NAR1 genes generally involve members of the formate/nitrite transporter family (Rexach et al., 2000). In fact, LCIA-expressing *Xenopus* oocytes display both low-affinity bicarbonate transport and high-affinity nitrite transport activities (Mariscal et al., 2006), suggesting that LCIA is involved in both bicarbonate uptake and nitrite uptake induced under low-CO₂ conditions. In other words, the suppression of LCIA by high-CO₂ conditions may reduce nitrogen availability in the chloroplast. Additionally, the molecular structure of the high-affinity-bicarbonate transporter *cmpABCD* is very similar to that of the nitrate/nitrite transporter *nrtABCD* in *Synechococcus* sp. PCC7942 (for review, see Badger & Price, 2003). The expression of high-affinity nitrate and nitrite transporter (HANT/HANiT) system IV is triggered by a sensing signal of low CO₂ but not NH₄⁺ (Galván et al., 1996; Rexach et al., 1999). These data suggest that changes in CO₂ concentration may also affect intracellular nitrogen availability. Further study should be conducted to identify the cooperative effect of CO₂ and nitrogen availability on the expression of CO₂, nitrogen, and gametogenesis-responsive proteins.




Fig. 4. Schematic illustration of a C/N-status model in low- (A) and high-CO₂-acclimated cells (B) under respective CO₂ conditions produced during acclimation in *C. reinhardtii*. Dissolved inorganic carbon and nitrogen species drawn in bold dominate. CA, carbonic anhydrases; CA2, CAH2 (Fujiwara et al., 1990; Rawat & Moroney, 1991; Tachiki et al., 1992); NiR, nitrite reductase; NR, nitrate reductase; PG, 2-phosphoglycolate; PGA, 3-phosphoglycerate; PSII, photosystem II; Rh, Rh1 (Soupeine et al., 2002; Yoshihara et al., 2008); T, (putative) transporters; Ta, LCIA (Duanmu et al., 2009; Mariscal et al., 2006); Tb, HANT/HANi system IV (Galván et al., 1996; Rexach et al., 1999). CCM models of WT/LC cells, inorganic nitrogen assimilation, and photorespiratory carbon oxidation in *C. reinhardtii* are modified from Yamano et al. (2010), Fernández et al. (2009), and Spalding (2009), respectively.

CAH2 was first reported as an active α -type carbonic anhydrase induced under high-CO₂ conditions and light (Fujiwara et al., 1990; Rawat & Moroney, 1991; Tachiki et al., 1992), but it is poorly expressed and located in the periplasmic space (Rawat & Moroney, 1991). However, the physiological roles and expressional regulation of high-CO₂-inducible CAH2 are not well understood. Another high-CO₂-inducible protein, Rh1, has been identified as a

human Rhesus protein in a homology search and is a paralog of the ammonium and/or CO₂ channels (Soupene et al., 2002). The lack of Rh1 impairs cell growth in *C. reinhardtii* under high-CO₂ conditions (Soupene et al., 2004). Fong et al. (2007) proposed that Rh proteins served as H₂CO₃ transporters in *Escherichia coli* under high-CO₂ conditions. Rh1 was originally expected to be located on the chloroplast envelope *in silico* but the Rh1-GFP fusion protein is located in the plasma membrane in transgenic *C. reinhardtii* cells (Yoshihara et al., 2008).

Some mechanisms of CCM, the photorespiratory nitrogen cycle, and the nitrate/nitrite transport system, and the interactions among them, are summarized in relation to high- and low-CO₂-acclimated cells in Figure 4.

6. High-CO₂ signaling

How can microalgal cells sense the CO₂ signal and respond to changes in CO₂ concentration? The most abundant extracellular carbonic anhydrase, CAH1, in low-CO₂ cells is replaced by high-CO₂-inducible extracellular 43 kDa protein/Fe-assimilation 1 (H43/FEA1) when low-CO₂-cells are transferred to high-CO₂ conditions (Allen et al., 2007; Baba et al., 2011a; Hanawa et al., 2004, 2007; Kobayashi et al., 1997). We found that H43/FEA1 was the most abundant extracellular soluble protein, which occupied about 26% of the total extracellular proteins of high (3%)-CO₂-grown cells for 3 days (Baba et al., 2011b). *H43/FEA1* homologous genes are found in the genomic sequences of the chlorophytes *Scenedesmus obliquus*, *Chlorococcum littorale*, and *Volvox carteri*, and the dinoflagellate *Heterocapsa triquetra* (Allen et al., 2007). This suggests that the *H43/FEA1* orthologs may be widely distributed among at least chlorophyte algae.

The function of H43/FEA1 is not completely understood but one possible role may be in iron assimilation (Allen et al., 2007; Rubinelli et al., 2002). Allen et al. (2007) identified *FEA1*, *FEA2*, and a candidate ferrireductase (*FRE1*) are expressed coordinately with iron assimilation components, and it was hypothesized that the proteins may facilitate iron uptake with high affinity by concentrating iron in the vicinity of the cells (Allen et al., 2007). *FEA1* and *FRE1* homologs were previously identified as the high-CO₂-responsive genes *HCR1* and *HCR2* in the marine chlorophyte *C. littorale*, suggesting that the components of the iron-assimilation pathway are responsive to changes in CO₂ concentration (Sasaki et al., 1998). A homology search of DNA sequences showed that *H43*, *FEA1*, and *HCR1* are identical (Allen et al., 2007; Hanawa et al., 2007), indicating that *H43/FEA1* expression was also induced by iron deficiency with transcriptional regulation. Therefore, we proposed that the gene is expressed as *H43/FEA1* (Baba et al., 2011a, 2011b).

In *C. reinhardtii*, 0.3% (v/v) CO₂ in air is sufficient to trigger the expression of the high-CO₂-inducible *H43/FEA1* and expression is correlated linearly between 0.04% and 0.3% (Hanawa et al., 2007). *H43/FEA1* can also be induced under heterotrophic conditions in the presence of acetate as an organic carbon source even under low-CO₂ conditions (Hanawa et al., 2007). In a previous study, the dCO₂ concentration in a cell suspension increased about 28 times from 1 to approximately 28 μM, which was identical to that equilibrated under the bubbling of 0.22% CO₂ in light, when cells were incubated in the presence of acetate and 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) (Hanawa et al., 2007). From these data, the authors concluded that the induction of *H43/FEA1* is triggered by the CO₂ signal, even CO₂ generated from respiration, but not acetate itself or the change in carbon metabolite

abundance. Thus, *H43/FEA1* expression can be regulated by a high-CO₂ signal at the transcriptional level, irrespective of high-CO₂ conditions. *H43/FEA1* is highly reliable as a high-CO₂ response marker. The signal for *H43/FEA1* expression might be sensed by putative proteins localized on the cell membrane, which are influenced by protein modifiers and send the signal for *H43/FEA1* expression (Hanawa et al., 2007).

H43/FEA1 expression is induced under excessive levels of Cd (>25 μM) or iron-deficient conditions (<1 μM) (Allen et al., 2007; Rubinelli et al., 2002). Fei et al. (2009) reported two transcriptional *cis*-elements that are responsive to the Fe-deficient signal (FeREs) for *H43/FEA1* expression, namely FeRE1 and FeRE2, which are located at -273/-259 and -106/-85 upstream from the *H43/FEA1* transcriptional initiation site. The conserved sequence motif was identified from some iron-deficiency-inducible genes (Fei et al., 2009). However, according to our recent study, the two *cis*-elements are not necessary for the high-CO₂-induced expression of the *H43/FEA1* gene (Baba et al., 2011a). The high-CO₂-responsive *cis*-element (HCRE) was located at a -537/-370 upstream region from the *H43/FEA1* transcriptional initiation site, although the precise location has not yet been determined (Baba et al., 2011a). These results show that *H43/FEA1* expression is regulated by the high-CO₂ signal alone via the HCRE, which is located distantly from the iron-deficient-responsive element. This observation indicates that *H43/FEA1* is a multi-signal-regulated gene (Fig. 5). We have not yet determined whether all of these signals may affect the expression of other high-CO₂-inducible proteins (Baba et al., 2011b). Allen et al. (2007) reported some proteins that are iron-deficient-responsive but not CO₂-responsive, so those proteins are considered components of the iron-assimilation system. In addition, an iron-assimilation component was not found among high-CO₂-inducible extracellular proteins analyzed experimentally (Baba et al., 2011b). The expression by either high-CO₂ or iron-deficient signals is a unique feature of *H43/FEA1*.

The regulation of CCM-related gene expression, which is positively induced by a low-CO₂ signal and negatively induced by a high-CO₂ signal, has been well characterized in *C. reinhardtii*. A zinc-finger protein named CCM1/CIA5 has been identified as a candidate of the CCM master regulator (Fukuzawa et al., 2001; Miura et al., 2004; Xiang et al., 2001). CCM1/CIA5 is a protein complex with a molecular mass of approximately 290–580 kDa that is induced independently by DIC availability; Zn is necessary for its enzymatic function (Kohinata et al., 2008). One of the CCM1/CIA5-mediated signaling systems functions in the expression of *CAH1*, which encodes a low-CO₂-inducible periplasmic carbonic anhydrase (Fukuzawa et al., 1990) and the signaling is mediated by a Myb-type transcriptional regulator named LCR1 (Yoshioka et al., 2004). CCM1/CIA5 may possibly function as an amplifier for the CO₂ signaling cascade (Yamano et al., 2008). A direct signaling factor for CCM induction has not been identified, although some candidates have been reported (Giordano et al., 2005; Kaplan & Reinhold, 1999; Yamano et al., 2008). The *CCM1/CIA5* mutant lacks suppression of *H43/FEA1* expression under both low-CO₂ and iron-sufficient conditions (Allen et al., 2007), suggesting that *H43/FEA1* expression is regulated by the CCM1/CIA5-dependent signaling cascade. However, the regulatory mechanism seems to be complex. The responses of *CAH1* and *H43/FEA1* expression are not an all-or-none type to the signals for a change in environmental CO₂ concentration, acetate concentration, and light intensity (Hanawa et al., 2007). Signaling for *H43/FEA1* expression may be partially associated with CCM1/CIA5 signaling, although additional signals may also exist (Fig. 5). *CAH2* is continuously expressed in a *CCM1/CIA5* mutant independent of CO₂ concentration

(Rawat & Moroney, 1991), suggesting that CAH2 expression is regulated by CCM1/CIA5 in the wild type. However, another high-CO₂-inducible protein, Rh1, is not likely regulated by CCM1/CIA5 (Wang et al., 2005).




Fig. 5. Schematic model of high-CO₂ signaling for *H43/FEA1* induction. Solid and broken lines are expected and putative signaling flows, respectively. Low-CO₂ signaling is modified from Miura et al. (2004) and Yamano and Fukuzawa (2009). Iron-deficient-inducible genes are according to Allen et al. (2007). Cd signaling on *H43/FEA1* induction, proposed by Rubinelli et al. (2002), is not drawn because little about it is known.

7. Conclusion

Compared to low-CO₂-inducible mechanisms that are well understood, analyses of high-CO₂-responsive mechanisms in microalgae at the molecular level have just started using the unicellular green alga *C. reinhardtii*. An accurate characterization of the acclimation mechanisms to high-CO₂ conditions will be important for both a detailed understanding of sensing and responding to environmental CO₂ changes and maximizing algal biomass productivity in mass cultivation. *H43/FEA1*, the most abundant extracellular protein in high-CO₂-acclimated cells, is expressed in response to multiple signals, including high-CO₂, iron-deficiency, or Cd-stress conditions. This suggests that, in addition to the high-CO₂ signal itself, abnormally stressful conditions such as strong nutrient depletion caused by rapid growth under high-CO₂ conditions may trigger expression of the gene. Targeted proteomics of whole *C. reinhardtii* established by Wienkoop et al. (2010) and a cDNA array (Yamano et al., 2008) or transcriptomics (Yamano & Fukuzawa, 2009), which has been applied to an expression analysis of CCM-associated genes, would be useful for further detailed analysis of high-CO₂ response phenomena. Our recent data indicate that the expression of gametogenesis-related proteins, which are strictly regulated by nitrogen availability, is triggered by high-CO₂ signals with a drastic change in extracellular proteins. These gametogenesis-related proteins in the periplasmic space of *C. reinhardtii* cells may play novel and crucial roles when *C. reinhardtii* is grown under high-CO₂ conditions.

8. Acknowledgments

This work was financially supported, in part, by a Grant-in-Aid for Scientific Research (Basic Research Area (S), No. 2221003 to YS) from the Japan Society for the Promotion of Science, the Core Research of Evolutional Science & Technology program (CREST) from the Japan Science and Technology Agency (JST) (to MB & YS), another fund from the Japan Science and Technology Agency (CREST/JST, to YS), and by the Global Environment Research Fund from the Japanese Ministry of Environment (FY2008-2010) to YS.

9. References

- Abe, J.; Kubo, T.; Takagi, Y.; Saito, T.; Miura, K.; Fukuzawa, H. & Matsuda, Y. (2004) The Transcriptional Program of Synchronous Gametogenesis in *Chlamydomonas reinhardtii*. *Current Genetics*, Vol.46, No.5, (November 2004), pp. 304–315, ISSN 0172-8083.
- Adair, W.S. (1985) Characterization of *Chlamydomonas* Sexual Agglutinins. *Journal of Cell Science*, Vol.2, Supplement, (1985), pp. 233–260, ISSN 0269-3518
- Aizawa, K. & Miyachi, S. (1986) Carbonic Anhydrase and CO₂ Concentrating Mechanisms in Microalgae and Cyanobacteria. *FEMS Microbiology Letters*, Vol.39, No.3, (August 1986), pp. 215–233, ISSN 0378-1097.
- Allen, M.D.; del Campo, J.A.; Kropat, J. & Merchant, S.S. (2007) *FEA1*, *FEA2*, and *FRE1*, Encoding Two Homologous Secreted Proteins and a Candidate Ferrireductase, Are Expressed Coordinately with *FOX1* and *FTR1* in Iron-Deficient *Chlamydomonas reinhardtii*. *Eukaryotic Cell*, Vol.6, No.10, (October 2007), pp. 1841–1852, ISSN 1535-9778.
- Baba, M.; Hanawa, Y.; Suzuki, I. & Shiraiwa, Y. (2011a) Regulation of the Expression of *H43/Fea1* by Multi-Signals. *Photosynthesis research*, Epub ahead of print, (January 2011), ISSN 1573-5079.
- Baba, M.; Suzuki, I. & Shiraiwa, Y. (2011b) Proteomic Analysis of High-CO₂-Inducible Extracellular Proteins in the Unicellular Green Alga, *Chlamydomonas reinhardtii*. *Plant & cell physiology*, Epub ahead of print, (June 2011), ISSN 0032-0781.
- Badger, M.R. & Price, G.D. (1994) The Role of Carbonic Anhydrase in Photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol.45, (June 1994), pp. 369–392, ISSN 1040-2519.
- Badger, M.R. & Price, G.D. (2003) CO₂ Concentrating Mechanisms in Cyanobacteria: Molecular Components, Their Diversity and Evolution. *Journal of experimental botany*, Vol.54, No.383, (February 2003), pp. 609–622, ISSN 0022-0957.
- Badger, M.R.; von Caemmerer, S.; Ruuska, S. & Nakano, H. (2000) Electron Flow to Oxygen in Higher Plants and Algae: Rates and Control of Direct Photoreduction (Mehler reaction) and Rubisco Oxygenase. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, Vol.355, No.1402, (October 2000), pp. 1433–1446, ISSN 0962-8436.
- Badger, M.R.; Price, G.D.; Long, B.M. & Woodger, F.J. (2006) The Environmental Plasticity and Ecological Genomics of the Cyanobacterial CO₂ Concentrating Mechanism. *Journal of experimental botany*, Vol.57, No.2, (2006), pp. 249–265, ISSN 0022-0957.
- Balkos, K.D. & Colman, B. (2007) Mechanism of CO₂ Acquisition in an Acid-Tolerant *Chlamydomonas*. *Plant, cell & environment*, Vol.30, No.6, (June 2007), pp. 745–752, ISSN 0140-7791.

- Bartlett, S.G.; Mitra, M. & Moroney, J.V. (2007) CO₂ Concentrating Mechanisms, In: *The Structure and Function of Plastids.*, Edited by Wise, R.R & Hooper J.K., pp. 253–271. Springer, ISBN 978-1-4020-4061-0, Dordrecht, The Netherlands.
- Beardall, J. & Giordano, M. (2002) Ecological Implications of Microalgal and Cyanobacterial CCMs and Their Regulation. *Functional Plant Biology*, Vol.29, No.2-3, (2002), pp. 335–347, ISSN 1445-4408.
- Bhatti, S. & Colman, B. (2008) Inorganic Carbon Acquisition by Some Synurophyte Algae. *Physiologia plantarum*, Vol.133, No.1, (February 2008), pp. 33–40, ISSN 0031-9317
- Box, G.E.P. & Wilson, K.B. (1951) On the Experimental Attainment of Optimum Conditions. *Journal of the Royal Statistical Society Series B*, Vol.13, No.1, (1951), pp. 1–45, ISSN 0035-9246.
- Buchanan, M.J. & Snell, W.J. (1988) Biochemical Studies on Lysin, a Cell Wall Degrading Enzyme Released During Fertilization in *Chlamydomonas*. *Experimental cell research*, Vol.179, No.1, (November 1988), pp. 181–193, ISSN 0014-4827.
- Buyanovsky, G.A. & Wagner, G.H. (1983) Annual Cycles of Carbon Dioxide Level in Soil Air. *Soil Science Society of America Journal*, Vol.47, No.6, (1983), pp. 1139–1145, ISSN 0361-5995.
- Chihara, M.; Nakayama, T.; Inouye, I. & Kodama, M. (1994) *Chlorococcum littorale*, a New Marine Green Coccoid Alga (Chlorococcales, Chlorophyceae). *Archiv für Protistenkunde*, Vol.144, No.3, (1994), pp. 227–235, ISSN 0003-9365.
- Collins, S. & Bell, G. (2004) Phenotypic Consequences of 1,000 Generations of Selection at Elevated CO₂ in a Green Alga. *Nature*, Vol. 431, No.7008, (September 2004), pp. 566–569, ISSN 0028-0836.
- Collins, S. & Bell, G. (2006) Evolution of Natural Algal Populations at Elevated CO₂. *Ecology letters*, Vol.9, No.2, (February 2006), pp. 129–135, ISSN 1461-023X.
- Collins, S.; Sültemeyer, D. & Bell, G. (2006) Changes in C Uptake in Populations of *Chlamydomonas reinhardtii* Selected at High CO₂. *Plant, cell & environment*, Vol.29, No.9, (September 2006), pp. 1812–1819, ISSN 0140-7791.
- Colman, B. & Balkos, K.D. (2005) Mechanisms of Inorganic Carbon Acquisition by *Euglena* species. *Canadian journal of botany. Journal canadien de botanique*, Vol.83, No.7, (July 2005), pp. 865–871, ISSN 0008-4026.
- Condie, K.C. & Sloan, R.E. (January 1998) *Origin and Evolution of Earth: Principles of Historical Geology*. Prentice-Hall, ISBN 978-0134918204, NJ.
- Demidov, E.; Iwasaki, I.; Satoh, A.; Kurano, N. & Miyachi, S. (2000) Short-term Responses of Photosynthetic Reactions to Extremely High-CO₂ Stress in a “High-CO₂” Tolerant Green Alga, *Chlorococcum littorale* and an Intolerant Green Alga *Stichococcus bacillaris*. *Russian journal of plant physiology : a comprehensive Russian journal on modern phytophysiology*, Vol.47, No.5, (September 2000), pp. 622–631, ISSN 1021-4437.
- Diaz, M.M. & Maberly, S.C. (2009) Carbon-Concentrating Mechanisms in Acidophilic Algae. *Phycologia*, Vol.48, No.2, (March 2009), pp. 77–85, ISSN 0031-8884.
- Doney, S.C.; Fabry, V.J.; Feely, R.A. & Kleypas, J.A. (2009) Ocean Acidification: the Other CO₂ Problem, *Annual review of marine science*, Vol.1, (January 2009), pp. 169–192, ISSN 1941-1405.
- Duanmu, D.; Miller, A.; Horken, K.; Weeks, D. & Spalding, M.H. (2009) Knockdown of Limiting-CO₂-Induced Gene *HLA3* Decreases HCO₃⁻ Transport and Photosynthetic Ci Affinity in *Chlamydomonas reinhardtii*. *Proceedings of the National Academy of*

- Sciences of the United States of America*, Vol.106, No.14, (April 2009), pp. 5990–5995, ISSN 0027-8424.
- Ertl, H.; Hallmann, A.; Wenzl, S. & Sumper, M. (1992) A Novel Extensin That May Organize Extracellular Matrix Biogenesis in *Volvox carteri*. *The EMBO journal*, Vol.11, No.6, (June 1992), pp. 2055–2062, ISSN 0261-4189.
- Falkowski, P.G. & Raven, J.A. (2007) *Aquatic Photosynthesis, 2nd Ed.* Princeton University Press, ISBN 978-0691115511, Princeton, NJ.
- Fei, X.; Eriksson, M.; Yang, J. & Deng, X. (2009) An Fe Deficiency Responsive Element with a Core Sequence of TGGCA Regulates the Expression of *Fea1* in *Chlamydomonas reinhardtii*. *Journal of biochemistry*, Vol.146, No.2, (August 2009), pp. 157–166, ISSN 0021-924X.
- Fernández, E.; Llamas, Á. & Galván, A. (2009) Nitrogen Assimilation and its Regulation, In: *The Chlamydomonas Sourcebook, 2nd Ed, Vol. 2.*, Edited by Stern D.B., pp. 69–113. Academic Press, ISBN 978-0-12-370875-5, San Diego, CA.
- Fong, R.N.; Kim, K.-S.; Yoshihara, C.; Inwood, W. & Kustu, S. (2007) The W148L Substitution in the *Escherichia coli* Ammonium Channel AmtB Increases Flux and Indicates That the Substrate Is an Ion. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.104, No.47, (November 2007), pp. 18706–18711, ISSN 0027-8424.
- Fujiwara, S.; Fukuzawa, H.; Tachiki, A. & Miyachi, S. (1990) Structure and Differential Expression of Two Genes Encoding Carbonic Anhydrase in *Chlamydomonas reinhardtii*. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.87, No.24, (December 1990), pp. 9779–9783, ISSN 0027-8424.
- Fukuda, S.; Suzuki, I.; Hama, T. & Shiraiwa, Y. (2011) Compensatory Response of the Unicellular-Calcifying Alga *Emiliania huxleyi* (Coccolithophorales, Haptophyta) to Ocean Acidification. *Journal of Oceanography*, Vol.67, No.1, (February 2011), pp. 17–25, ISSN 0916-8370.
- Fukuzawa, H.; Fujiwara, S.; Yamamoto, Y.; Dionisio-Sese, M.L. & Miyachi, S. (1990) cDNA Cloning, Sequence, and Expression of Carbonic Anhydrase in *Chlamydomonas reinhardtii*: Regulation by Environmental CO₂ Concentration. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.87, No.11, (June 1990), pp. 4383–4387, ISSN 0027-8424.
- Fukuzawa, H.; Miura, K.; Ishizaki, K.; Kucho, K.I.; Saito, T.; Kohinata, T. & Ohyama, K. (2001) *Ccm1*, a Regulatory Gene Controlling the Induction of a Carbon-Concentrating Mechanism in *Chlamydomonas reinhardtii* by Sensing CO₂. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.98, No.9, (April 2001), pp. 5347–5352, ISSN 0027-8424.
- Galvan, A.; Quesada, A. & Fernández, E. (1996) Nitrate and Nitrite Are Transported by Different Specific Transport Systems and by a Bispecific Transporter in *Chlamydomonas reinhardtii*. *The Journal of biological chemistry*, Vol.271, No. 4, (January 1996), pp. 2088–2092, ISSN 0021-9258.
- Giordano, M.; Beardall, J. & Raven, J.A. (2005) CO₂ Concentrating Mechanisms in Algae: Mechanisms, Environmental Modulation, and Evolution. *Annual review of plant biology*, Vol.56, (June 2005), pp. 99–131, ISSN 1543-5008.
- Giordano, M.; Norici, A.; Forssen, M.; Eriksson, M. & Raven, J.A. (2003) An Anaplerotic Role for Mitochondrial Carbonic Anhydrase in *Chlamydomonas reinhardtii*. *Plant Physiology*, Vol.132, No.4, (August 2003), pp. 2126–2134, ISSN 0032-0889.

- Goodenough, U.W.; Gebhart, B.; Mecham, R.E. & Heuser, J.E. (1986) Crystals of the *Chlamydomonas reinhardtii* Cell Wall: Polymerization, Depolymerization, and Purification of Glycoprotein Monomers. *The Journal of cell biology*, Vol.103, No.2, (August 1986), pp. 405–417, ISSN 0021-9525.
- Goodenough, U.; Lin, H. & Lee, J. (2007) Sex Determination in *Chlamydomonas*. *Seminars in cell & developmental biology*, Vol.18, No.3, (February 2007), pp. 350–361, ISSN 1084-9521.
- Hallmann, A. (2006) The Pherophorins: Common, Versatile Building Blocks in the Evolution of Extracellular Matrix Architecture in *Volvocales*. *The Plant journal : for cell and molecular biology*, Vol.45, No.2, (January 2006), pp. 292–307, ISSN 0960-7412.
- Hanawa, Y. (2007) Study on a CO₂ Sensing Mechanism by the Expression Analysis of a High CO₂ Inducible H43 Gene in *Chlamydomonas reinhardtii*. A Dissertation Submitted to the Graduate School of Life and Environmental Sciences, the University of Tsukuba.
- Hanawa, Y.; Iwamoto, K. & Shiraiwa, Y. (2004) Purification of a Recombinant H43, a High-CO₂-Inducible Protein of *Chlamydomonas reinhardtii*, Expressed in *Escherichia coli*. *Japanese Journal of Phycology*, Vol.52, Supplement, (2004), pp. 95–100, ISSN 0038-1578.
- Hanawa, Y.; Watanabe, M.; Karatsu, Y.; Fukuzawa, H. & Shiraiwa, Y. (2007) Induction of a High-CO₂-Inducible, Periplasmic Protein, H43, and Its Application as a High-CO₂-Responsive Marker for Study of the High-CO₂-Sensing Mechanism in *Chlamydomonas reinhardtii*. *Plant & cell physiology*, Vol.48, No.2, (January 2007), pp. 299–309, ISSN 0032-0781.
- Heldt, H.; Werdan, K.; Milovancev, M. & Geller, G. (1973) Alkalization of the Chloroplast Stroma Caused by Light-Dependent Proton Flux into the Thylakoid Space. *Biochimica et Biophysica Acta*, Vol.314, No.2, (August 1973), pp. 224–241, ISSN 0006-3002.
- Ho, S.H.; Chen, C.Y.; Lee, D.J. & Chang, J.S. (2011) Perspectives on Microalgal CO₂-Emission Mitigation Systems - A Review. *Biotechnology Advances*, Vol.29, No.2, (March-April 2011), pp. 189–198, ISSN 0734-9750.
- Hoffmann, X.K. & Beck, C.F. (2005) Mating-Induced Shedding of Cell Walls, Removal of Walls from Vegetative Cells, and Osmotic Stress Induce Presumed Cell Wall Genes in *Chlamydomonas*. *Plant Physiology*, Vol.139, No.2, (October 2005), pp. 999–1014, ISSN 0032-0889.
- Hurd, C.L.; Hepburn, C.D.; Currie, K.I.; Raven, J.A. & Hunter, K.A. (2009) Testing the Effects of Ocean Acidification on Algal Metabolism: Considerations for Experimental Designs. *Journal of Phycology*, Vol.45, No.6, (November 2009), pp. 1236–1251, ISSN 0022-3646.
- Iglesias-Rodriguez, M.D.; Halloran, P.R.; Rickaby, R.E.M.; Hall, I.R.; Colmenero-Hidalgo, E.; Gittins, J.R.; Green, D.R.H.; Tyrrell, T.; Gibbs, S.J.; Dassow, P.; Rehm, E.; Armbrust, E.V. & Boessenkool, K.P. (2008) Phytoplankton Calcification in a High-CO₂ World. *Science*, Vol.320, No.5874, (April 2008), pp. 336–340, ISSN 0036-8075.
- Inoue, I. (2007) *The Natural History of Algae: Second Edition: Perspective of Three Billion years Evolution of Algae, Earth and Environment*. Tokai University Press, ISBN 978-4-486-01777-6, Japan.
- Ishii, M.; Yoshikawa, H. & Matsueda, H. (March, 2000) *Coulometric Precise Analysis of Total Inorganic Carbon in Seawater and Measurements of Radiocarbon for the Carbon Dioxide in*


- the Atmosphere and for the Total Inorganic Carbon in Seawater. Technical Reports of the Meteorological Research Institute No. 41.*, Meteorological Research Institute, Japan, Retrieved from <http://www.mri-jma.go.jp/Publish/Technical/index_en.html>
- Iwasaki, I.; Hu, Q.; Kurano, N. & Miyachi, S. (1998) Effect of Extremely High-CO₂ Stress on Energy Distribution Between Photosystem I and Photosystem II in a High-CO₂ Tolerant Green Alga, *Chlorococcum littorale* and the Intolerant Green Alga *Stichococcus bacillaris*. *Journal of photochemistry and photobiology. B, Biology*, Vol.44, No.3, (July 1998), pp. 184–190, ISSN 1011-1344.
- Iwasaki, I.; Kurano, N. & Miyachi, S. (1996) Effects of High-CO₂ Stress on Photosystem II in a Green Alga, *Chlorococcum littorale*, Which Has a Tolerance to High CO₂. *Journal of photochemistry and photobiology. B, Biology*, Vol.36, No.3, (December 1996), pp. 327–332, ISSN 1011-1344.
- Jones, H.G. (1992) *Plants and Microclimate: A Quantitative Approach to Environmental Plant Physiology, 2nd edition.*, Cambridge University Press, ISBN 9780521425247, Cambridge, UK.
- Kaplan, A. & Reinhold, L. (1999) CO₂ Concentrating Mechanisms in Photosynthetic Microorganisms. *Annual Review of Plant Physiology and Plant Molecular Biology*, Vol.50, (June 1999), pp. 539–570, ISSN 1040-2519.
- Kinoshita, T.; Fukuzawa, H.; Shimada, T.; Saito, T. & Matsuda, Y. (1992) Primary Structure and Expression of a Gamete Lytic Enzyme in *Chlamydomonas reinhardtii*: Similarity of Functional Domains to Matrix Metalloproteases. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.89, No.10, (May 1992), pp. 4693–4697, ISSN 0027-8424.
- Kobayashi, H.; Odani, S. & Shiraiwa, Y. (1997) A High-CO₂-Inducible, Periplasmic Polypeptide in an Unicellular Green Alga *Chlamydomonas reinhardtii* (abstract no. 493). *Plant Physiology*, Vol.114, Supplement, p. 112, ISSN 0032-0889.
- Kodama, M.; Ikemoto, H. & Miyachi, S. (1993) A New Species of Highly CO₂-Tolerant Fast-Growing Marine Microalga Suitable for High-Density Culture. *Journal of Marine Biotechnology*, Vol. 1, (1993), pp. 21–25, ISSN 0941-2905.
- Kohinata, T.; Nishino, H. & Fukuzawa, H. (2008) Significance of Zinc in a Regulatory Protein, CCM1, Which Regulates the Carbon-Concentrating Mechanism in *Chlamydomonas reinhardtii*. *Plant & cell physiology*, Vol.49, No.2, (January 2008), pp. 273–283, ISSN 0032-0781.
- Kubo, T.; Saito, T.; Fukuzawa, H. & Matsuda, Y. (2001) Two Tandemly-Located Matrix Metalloprotease Genes with Different Expression Patterns in the *Chlamydomonas* Sexual Cell Cycle. *Current Genetics*, Vol.40, No.2, (September 2001), pp. 136–143, ISSN 0172-8083.
- Kumar, A.; Ergas, S.J.; Yuan, X.; Sahu, A.K.; Zhang, Q.; Dewulf, J.; Malcata, F.X. & an Langenhove, H.V. (2010) Enhanced CO₂ Fixation and Biofuel Production via Microalgae: Recent Developments and Future Directions, *Trends in biotechnology*, Vol.28, No.7, (June 2010), pp. 371–380, ISSN 0167-7799.
- Kurano, N.; Ikemoto, H.; Miyashita, H.; Hasegawa, T.; Hata, H. & Miyachi, S. (1995) Fixation and Utilization of Carbon Dioxide by Microalgal Photosynthesis. *Energy Conversion and Management*, Vol.36, No.6–9, (June-September 1995), pp. 689–692, ISSN 0196-8904.

- Lee, J.S. & Lee, J.P. (2003) Review of Advances in Biological CO₂ Mitigation Technology. *Biotechnology and Bioprocess Engineering*, Vol.8, No.6, (2003), pp. 354–359, ISSN 1226-8372.
- Lee, J.H.; Waffenschmidt, S.; Small, L. & Goodenough, U. (2007) Between-Species Analysis of Short-Repeat Modules in Cell Wall and Sex-Related Hydroxyproline-Rich Glycoproteins of *Chlamydomonas*. *Plant Physiology*, Vol.144, No.4, (August 2007), pp. 1813–1826, ISSN 0032-0889.
- Maeda, K.; Owada, M.; Kimura, N.; Omata, K. & Karube, I. (1995) CO₂ Fixation from the Flue Gas on Coal-Fired Thermal Power plant by Microalgae. *Energy Conversion and Management*, Vol.36, No.6–9, (June–September 1995), pp. 717–720, ISSN 0196-8904.
- Mariscal, V.; Moulin, P.; Orsel, M.; Miller, A.J.; Emilio Fernández, E. & Galvána, A. (2006) Differential Regulation of the *Chlamydomonas* *Nar1* Gene Family by Carbon and Nitrogen. *Protist*, Vol.157, No.4, (October 2006), pp. 421–433, ISSN 1434-4610.
- Matsuda, Y.; Shimada, T. & Sakamoto, Y. (1992) Ammonium Ions Control Gametic Differentiation and Dedifferentiation in *Chlamydomonas reinhardtii*. *Plant & cell physiology*, Vol.33, No.7, (1992), pp. 909–914, ISSN 0032-0781.
- Miura, K.; Yamano, T.; Yoshioka, S.; Kohinata, T.; Inoue, Y.; Taniguchi, F.; Asamizu, E.; Nakamura, Y.; Tabata, S.; Yamato, K.T.; Ohyama, K. & Fukuzawa, H. (2004) Expression Profiling-Based Identification of CO₂-Responsive Genes Regulated by CCM1 Controlling a Carbon Concentrating Mechanism in *Chlamydomonas reinhardtii*. *Plant Physiology*, Vol.135, No.3, (July 2004), pp. 1595–1607, ISSN 0032-0889.
- Moroney, J.V. & Ynalvez, R.A. (2007) A Proposed Carbon Dioxide Concentration Mechanism in *Chlamydomonas reinhardtii*. *Eukaryotic Cell*, Vol.6, No. 8, (August 2007), pp. 1251–1259, ISSN 1535-9778.
- Nielsen, E.S. (1955) Carbon Dioxide As Carbon Source and Narcotic in Photosynthesis and Growth of *Chlorella pyrenoidosa*. *Physiologia Plantarum*, Vol.8, No.2, (April 1955), pp. 317–335, ISSN 0031-9317.
- Olaizola, M. (2003) Microalgal Removal of CO₂ from Flue Gases: Changes in Medium pH and Flue Gas Composition Do not Appear to Affect the Photochemical Yield of Microalgal Cultures. *Biotechnology and Bioprocess Engineering*, Vol.8, No.6, (November–December 2003), pp. 360–367, ISSN 1226-8372.
- Pesheva, I.; Kodama, M.; Dionisio-Sese, M.L. & Miyachi, S. (1994) Changes in Photosynthetic Characteristics Induced by Transferring Air-Grown Cells of *Chlorococcum littorale* to High-CO₂ Conditions. *Plant & cell physiology*, Vol.35, No.3, (1994), pp. 379–387, ISSN 0032-0781.
- Price, G.D. & Badger, M.R. (1989) Isolation and Characterization of High CO₂-Requiring-Mutants of the Cyanobacterium *Synechococcus* PCC7942, Two Phenotypes that Accumulate Inorganic Carbon but Are Apparently Unable to Generate CO₂ within the Carboxysome. *Plant Physiology*, Vol.91, No.2, (October 1989), pp. 514–525, ISSN 0032-0889.
- Pronina, N.A.; Kodama, M. & Miyachi, S. (1993) Changes in Intracellular pH Values in Various Microalgae Induced by Raising CO₂ Concentrations. *XV International Botanical Congress*, ISBN 9783921800386, Yokohama, Japan, August, 1993.
- Ramazanov, Z. & Cárdenas, J. (1994) Photorespiratory Ammonium Assimilation in Chloroplasts of *Chlamydomonas reinhardtii*. *Physiologia Plantarum*, Vol.91, No.3, (July 1994), pp. 495–502, ISSN 0031-9317.

- Raven, J.A. (2001) A Role for Mitochondrial Carbonic Anhydrase in Limiting CO₂ Leakage from Low CO₂-Grown Cells of *Chlamydomonas reinhardtii*. *Plant, Cell & Environment*, Vol.24, No.2, (February 2001), pp. 261–265, ISSN 0140-7791.
- Raven, J.A. (2010) Inorganic Carbon Acquisition by Eukaryotic Algae: Four Current Questions. *Photosynthesis research*, Vol.106, No.1-2, (June 2010), pp. 123–134, ISSN 1573-5079.
- Raven, J.A.; Cockell, C.S. & De La Rocha, C.L. (2008) The Evolution of Inorganic Carbon Concentrating Mechanisms in Photosynthesis. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, Vol.363, No.1504, (August 2008), pp. 2641–2650, ISSN 0962-8436.
- Rawat, M. & Moroney, J.V. (1991) Partial Characterization of a New Isoenzyme of Carbonic-anhydrase Isolated from *Chlamydomonas reinhardtii*. *The Journal of biological chemistry*, Vol.266, No.15, (May 1991), pp. 9719–9723, ISSN 0021-9258.
- Rexach, J.; Fernández, E. & Galván, A. (2000) The *Chlamydomonas reinhardtii* *Nar1* Gene Encodes a Chloroplast Membrane Protein Involved in Nitrite Transport. *The Plant Cell*, Vol.12, No.8, (August 2000), pp. 1441–1453, ISSN 1040-4651.
- Rexach, J.; Montero, B.; Fernández, E. & Galván, A. (1999) Differential Regulation of the High Affinity Nitrite Transport Systems III and IV in *Chlamydomonas reinhardtii*. *The Journal of biological chemistry*, Vol.274, No.39, (September 1999), pp. 27801–27806, ISSN 0021-9258.
- Riebesell, U.; Zondervan, I.; Rost, B.; Tortell, P.D.; Zeebe, R.E. & Morel, F.M.M. (2000) Reduced Calcification of Marine Plankton in Response to Increased Atmospheric CO₂. *Nature*, Vol.407, No.6802, (September 2000), pp. 364–367, ISSN 0028-0836.
- Rubinelli, P.; Siripornadulsil, S.; Gao-Rubinelli, F. & Sayre, R.T. (2002) Cadmium- and Iron-Stress-Inducible Gene Expression in the Green Alga *Chlamydomonas reinhardtii*: Evidence for H43 Protein Function in Iron Assimilation. *Planta*, Vol.215, No.1, (May 2002), pp. 1–13, ISSN 0032-0935.
- Sasaki, T.; Kurano, N. & Miyachi, S. (1998) Induction of Ferric Reductase Activity and of Iron Uptake Capacity in *Chlorococcum littorale* Cells under Extremely High-CO₂ and Iron-Deficient Conditions. *Plant & cell physiology*, Vol.39, No.4, (1998), pp. 405–410, ISSN 0032-0781.
- Sasaki, T.; Pronina, N.A.; Maeshima, M.; Iwasaki, I.; Kurano, N. & Miyachi, S. (1999) Development of Vacuoles and Vacuolar ATPase Activity under Extremely High-CO₂ Conditions in *Chlorococcum littorale* Cells. *Plant Biology*, Vol.1, No.1, (January 1999), pp. 68–75, ISSN 1435-8603.
- Satoh, A.; Kurano, N.; Harayama, S. & Miyachi, S. (2004) Effects of Chloramphenicol on Photosynthesis, Protein Profiles and Transketolase Activity under Extremely High CO₂ Concentration in an Extremely-high-CO₂-tolerant Green Microalga, *Chlorococcum littorale*. *Plant & Cell Physiology*, Vol.45, No.12, (December 2004), pp. 1857–1862, ISSN 0032-0781.
- Satoh, A.; Kurano, N. & Miyachi, S. (2001) Inhibition of Photosynthesis by Intracellular Carbonic Anhydrase in Microalgae under Excess Concentrations of CO₂. *Photosynthesis research*, Vol.68, No.3, (2001), pp. 215–224, ISSN 1573-5079.
- Satoh, A.; Kurano, N.; Senger, H. & Miyachi, S. (2002) Regulation of Energy Balance in Photosynthesis in Response to Changes in CO₂ Concentrations and Light Intensities During Growth in Extremely High-CO₂-Tolerant Green Microalgae. *Plant & cell physiology*, Vol.43, No.4, (April 2002), pp. 440–451, ISSN 0032-0781.

- Seckbach, J. Baker, F.A. & Shugarman, P.M. (1970) Algae Thrive under Pure CO₂. *Nature*, Vol.227, No.5259, (1970), pp. 744–745, ISSN 0028-0836.
- Shiraiwa, Y. & Miyachi, S. (1985) Effect of Temperature and CO₂ Concentration on Induction of Carbonic Anhydrase and Changes in Efficiency of Photosynthesis in *Chlorella vulgaris* 11h. *Plant & cell physiology*, Vol.26, No.3, (1985), pp. 543–549, ISSN 0032-0781.
- Shiraiwa, Y. & Schmid, G.H. (1986) Stimulation of photorespiration by the carbonic anhydrase inhibitor ethoxycarbonylamine in *Chlorella vulgaris*. *Zeitschrift für Naturforschung* Vol.41, No.5-6, (1986), pp.564–570, ISSN 0341-0382.
- Soupene, E.; Inwood, W. & Kustu, S. (2004) Lack of the Rhesus Protein Rh1 Impairs Growth of the Green Alga *Chlamydomonas reinhardtii* at High CO₂. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.101, No.20, (May 2004) :7787–7792, ISSN 0027-8424.
- Soupene, E.; King, N.; Feild, E.; Liu, P.; Niyogi, K.K.; Huang, C.-H. & Kustu, S. (2002) Rhesus Expression in Green Alga Is Regulated by CO₂. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.99, No.11, (May 2002), pp. 7769–7773, ISSN 0027-8424.
- Spalding, M.H. (2008) Microalgal Carbon-Dioxide-Concentrating Mechanisms: *Chlamydomonas* Inorganic Carbon Transporters. *Journal of experimental botany*, Vol.59, No.7, (2008), pp. 1463–1473, ISSN 0022-0957.
- Spalding, M.H. (2009) The CO₂-Concentrating Mechanism and Carbon Assimilation, In: *The Chlamydomonas Sourcebook, 2nd Ed, Vol. 2.*, Edited by Stern D.B., pp. 257–301. Academic Press, ISBN 978-0-12-370875-5, San Diego, CA.
- Spalding M.H.; Spreitzer, R.J. & Ogren, W.L. (1983) Reduced Inorganic Carbon Transport in a CO₂-Requiring Mutant of *Chlamydomonas reinhardtii*. *Plant Physiology*, Vol.73, No.2, (October 1983), pp. 273–276, ISSN 0032-0889.
- Stolzy, L.H. (1974) Soil Atmosphere. In: *The Plant Root and Its Environment.*, Edited by Carson E.W., pp 335–362. University Press of Virginia, ISBN 978-0813904115, Charlottesville, VA.
- Suzuki, K. & Spalding, M.H. (1989) Adaptation of *Chlamydomonas reinhardtii* High-CO₂-Requiring Mutants to Limiting CO₂. *Plant Physiology*, Vol.90, No.3, (July 1989), pp. 1195–1200, ISSN 0032-0889.
- Sültemeyer, D.F.; Klug, K. & Fock, H.P. (1986) Effect of Photon Fluence Rate on Oxygen Evolution and Uptake by *Chlamydomonas reinhardtii* Suspensions Grown in Ambient and CO₂-Enriched Air. *Plant Physiology*, Vol.81, No.2, (June 1986), pp. 372–375, ISSN 0032-0889.
- Sültemeyer, D.F.; Klug, K. & Fock, H.P. (1987) Effect of Dissolved Inorganic Carbon on Oxygen Evolution and Uptake by *Chlamydomonas reinhardtii* Suspensions Adapted to Ambient and CO₂-Enriched Air. *Photosynthesis research*, Vol.12, No.1, (1987), pp. 25–33, ISSN 1573-5079.
- Tachiki, A.; Fukuzawa, H. & Miyachi, S. (1992) Characterization of Carbonic-anhydrase Isozyme CA2, Which Is the CAH2 Gene-product, in *Chlamydomonas reinhardtii*. *Bioscience, Biotechnology, & Biochemistry*, Vol.56, No.5, (May 1992), pp. 794–798, ISSN 0916-8451.
- Verma, V.; Bhati, S.; Huss, V.A.R. & Colman, B. (2009) Photosynthetic Inorganic Carbon Assimilation in a Free-Living Species of *Coccomyxa* (Chlorophyta). *Journal of Phycology*, Vol.45, (2009), pp. 847–854, ISSN 0022-3646.

- Wang, Y.; Sun, Z.; Horken, K.M.; Im, C.S.; Xiang, Y.; Grossman, A.R. & Weeks, D.P. (2005) Analysis of CIA5, the Master Regulator of the Carbon-concentrating Mechanisms in *Chlamydomonas reinhardtii*, and Its Control of Gene Expression. *Canadian Journal of Botany*, Vol.83, No.7, (July 2005), pp. 765–779, ISSN 0008-4026.
- Wienkoop, S.; Weiss, J.; May, P.; Kempa, S.; Irgang, S.; Recuenco-Munoz, L.; Pietzke, M.; Schwemmer, T.; Rupprecht, J.; Egelhofer, V. & Weckwerth, W. (2010) Targeted Proteomics for *Chlamydomonas reinhardtii* Combined with Rapid Subcellular Protein Fractionation, Metabolomics and Metabolic Flux Analyses. *Molecular BioSystems*, Vol.6, No. 6, (June 2010), pp. 1018–1031, ISSN 1742-2051.
- Xiang, Y.; Zhang, J. & Weeks, D.P. (2001) The *Cia5* Gene Controls Formation of the Carbon Concentrating Mechanism in *Chlamydomonas reinhardtii*. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.98, No. 9, (April 2001), 5341–5346, ISSN 0027-8424.
- Yamano, T. & Fukuzawa, H. (2009) Carbon-Concentrating Mechanism in a Green Alga, *Chlamydomonas reinhardtii*, Revealed by Transcriptome Analyses. *Journal of basic microbiology*, Vol. 49, No.1, (February 2009), pp. 42–51, ISSN 0233-111X.
- Yamano, T.; Miura, K. & Fukuzawa, H. (2008) Expression Analysis of Genes Associated with the Induction of the Carbon-Concentrating Mechanism in *Chlamydomonas reinhardtii*. *Plant Physiology*, Vol.147, No.1, (May 2008), pp. 340–354, ISSN 0032-0889.
- Yamano, T.; Tsujikawa, T.; Hatano, K.; Ozawa, S.; Takahashi, Y. & Fukuzawa, H. (2010) Light and Low-CO₂-Dependent LCIB-LCIC Complex Localization in the Chloroplast Supports the Carbon Concentrating Mechanism in *Chlamydomonas reinhardtii*. *Plant & cell physiology*, Vol.51, No.9, (July 2010), pp. 1453–1468, ISSN 0032-0781.
- Yang, S.Y.; Tsuzuki, M. & Miyachi, S. (1985) Carbonic Anhydrase of *Chlamydomonas*: Purification and Studies on Its Induction Using Antiserum Against *Chlamydomonas* Carbonic Anhydrase. *Plant & cell Physiology*, Vol.26, No.1, (1985), pp. 25–34, ISSN 0032-0781.
- Yoshihara, C.; Inoue, K.; Schichnes, D.; Ruzin S.; Inwood, W. & Kustu, S. (2008) An Rh1-GFP Fusion Protein Is in the Cytoplasmic Membrane of a White Mutant Strain of *Chlamydomonas reinhardtii*. *Molecular Plant*, Vol.1, No.6, (November 2008), pp. 1007–1020, ISSN 1674-2052.
- Yoshioka, S.; Taniguchi, F.; Miura, K.; Inoue, T.; Yamano, T. & Fukuzawa, H. (2004) The Novel Myb Transcription Factor LCR1 Regulates the CO₂-Responsive Gene *Cah1*, Encoding a Periplasmic Carbonic Anhydrase in *Chlamydomonas reinhardtii*. *The Plant Cell*, Vol.16, No.6, (June 2004), pp. 1466–1477, ISSN 1040-4651.
- Yun, Y.S. & Park, J.M. (1997) Development of Gas Recycling Photobioreactor System for Microalgal Carbon Dioxide Fixation. *Korean Journal of Chemical Engineering*, Vol.14, No.4, (1997), pp. 297–300, ISSN 0256-1115.



Advances in Photosynthesis - Fundamental Aspects

Edited by Dr Mohammad Najafpour

ISBN 978-953-307-928-8

Hard cover, 588 pages

Publisher InTech

Published online 15, February, 2012

Published in print edition February, 2012

Photosynthesis is one of the most important reactions on Earth. It is a scientific field that is the topic of many research groups. This book is aimed at providing the fundamental aspects of photosynthesis, and the results collected from different research groups. There are three sections in this book: light and photosynthesis, the path of carbon in photosynthesis, and special topics in photosynthesis. In each section important topics in the subject are discussed and (or) reviewed by experts in each book chapter.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Masato Baba and Yoshihiro Shiraiwa (2012). High-CO₂ Response Mechanisms in Microalgae, *Advances in Photosynthesis - Fundamental Aspects*, Dr Mohammad Najafpour (Ed.), ISBN: 978-953-307-928-8, InTech, Available from: <http://www.intechopen.com/books/advances-in-photosynthesis-fundamental-aspects/high-co2-response-mechanism-in-microalgae>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.